



2 Jungle Road, Leominster, MA. 01453 | 386 Elm Street, Gardner, MA. 01440

Tel: 978-534-8300 Fax: 978-840-8508 | Tel: 978-632-7270 Fax: 978-632-7198

[www.centralmassoms.com](http://www.centralmassoms.com)

Carmine A. Colarusso, D.M.D.

James F. Connors II, D.M.D., M.D.

Shant J. Baran, D.M.D.

## CONSENT FOR INTRAVENOUS SEDATION/ANESTHESIA

---

Patient's Name

Date

**Please read each paragraph carefully. If you have any questions about your proposed treatment, please ask your doctor BEFORE signing.**

You have chosen **intravenous sedation** for your surgery, a common procedure that is considered quite safe. Nevertheless, any anesthesia carries some risk and the common ones known for intravenous anesthesia are noted below for your review before you consent to its use:

- Allergic reactions (previously unknown) to any of the medications used.
- Discomfort, swelling or bruising at the site where the drugs are placed into a vein.
- Vein irritation, called phlebitis, where the needle is placed into a vein. Sometimes this may progress to a level of discomfort where arm or hand motion may be restricted, or further medication or care may be required.
- Nausea and vomiting, although not common, are unfortunate side effects of intravenous anesthesia. Bed rest, and sometimes medications, may be required for relief.
- Intravenous sedation is a serious medical procedure and, whether given in a hospital or office, carries with it the risk of brain damage, stroke, heart attack or death.

### YOUR OBLIGATIONS:

- Because the anesthetic medication (including oral premedication/sedation) causes prolonged drowsiness, you **MUST** be accompanied by a responsible adult to drive you to and from surgery, and stay with you for several hours until you are recovered sufficiently to care for yourself. Sometimes the effects of the drugs do not wear off for 24 hours.
- During recovery time (normally 24 hours), you should not drive, operate complicated machinery or devices or make important decisions such as signing documents, etc.
- You must have a completely empty stomach. It is vital that you have **NO SOLID FOODS OR DAIRY PRODUCTS six (6) hours** prior to your anesthetic AND **NOTHING AT ALL TO EAT OR DRINK INCLUDING LIQUIDS FOR two (2) hours** prior to your anesthetic. **TO DO OTHERWISE MAY BE LIFE-THREATENING!**

### CONSENT

I have read and understand the above paragraphs and realize that intravenous sedation carries with it certain serious risks. I request that intravenous anesthesia be used for my surgery. All my questions

regarding this consent have been answered fully and to my satisfaction, and I fully understand the risks involved. I certify that I speak, read and write English.

---

Patient's (or Legal Guardian's) Signature

Date

---

Doctor's Signature

Witness' Signature

Date

Rev 9/2010